

Janusz Gajda
 Institute of Mathematics and Computer Science
 Wrocław University of Technology
 E-mail: janusz.gajda@pwr.wroc.pl

Financial applications of subordinated tempered α -stable processes

We propose a new model (see [1]) for financial data description. This model is a combination of tempered stable process $T(\tau)$ [2, 3] and inverse stable subordinator $S(t)$ [4]. As a result we obtain process $X(t) = T(S(t))$. We classify our system to the family of subdiffusive processes and investigate its tail behavior. Moreover we propose a Fokker-Planck equation (FPE) for the density $f(x, t)$ of the process $X(t)$

$$\frac{\partial f(x, t)}{\partial t} = {}_0D_t^{1-\gamma} \frac{\partial^{\alpha, \lambda}}{\partial x} f(x, t). \quad (1)$$

This FPE consists of two fractional operators. First one is the time operator of fractional Riemann-Liouville [5] derivative connected with inverse stable subordinator. The second one is the spatial operator connected with the external tempered stable process [3].

In the last step we calibrate our model to the real data. The process $X(t)$ captures not only the tempered stable character of the underlying data but also such property as periods in which the values of an asset stay on the same level. In addition we describe in details testing and estimation procedures.

References

- [1] J. Gajda, A. Wyłomańska, *Physica A* 392 (2013), 3168.
- [2] J. Rosiński, *Stochastic Processes and Their Applications* 117 (2007), 677.
- [3] B. Baeumer, M. M. Meerschaert, *J. Comput. Appl. Math.* 233 (2010), 2438.
- [4] M. Magdziarz, A. Weron, K. Weron, *Phys. Rev. E* 75 (2007), 016708.
- [5] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies 204, Elsevier, 2006.
- [6] G. Samorodnitsky, M. S. Taqqu, *Stable Non-Gaussian Random Processes: Stochastic Models with Infinite Variance*, Chapman & Hall, London, 1994.