



University at Buffalo
The State University of New York

**THERE AND BACK AGAIN:
FROM THE BORSUK-ULAM THEOREM
TO QUANTUM SPACES**

Piotr M. Hajac (IMPAN)

Joint work with
P. F. Baum, L. Dąbrowski, S. Neshveyev and T. Maszczyk.

Jiří Matoušek

Using the Borsuk-Ulam Theorem

Lectures on Topological Methods
in Combinatorics and Geometry



The Borsuk-Ulam Theorem

Theorem (Borsuk-Ulam)

Let n be a positive natural number. If $f: S^n \rightarrow \mathbb{R}^n$ is continuous, then there exists a pair $(p, -p)$ of antipodal points on S^n such that $f(p) = f(-p)$.

The Borsuk-Ulam Theorem

Theorem (Borsuk-Ulam)

Let n be a positive natural number. If $f: S^n \rightarrow \mathbb{R}^n$ is continuous, then there exists a pair $(p, -p)$ of antipodal points on S^n such that $f(p) = f(-p)$.

The logical negation of the theorem

There exists a continuous map $f: S^n \rightarrow \mathbb{R}^n$ such that for all pairs $(p, -p)$ of antipodal points on S^n we have $f(p) \neq f(-p)$.

The Borsuk-Ulam Theorem

Theorem (Borsuk-Ulam)

Let n be a positive natural number. If $f: S^n \rightarrow \mathbb{R}^n$ is continuous, then there exists a pair $(p, -p)$ of antipodal points on S^n such that $f(p) = f(-p)$.

The logical negation of the theorem

There exists a continuous map $f: S^n \rightarrow \mathbb{R}^n$ such that for all pairs $(p, -p)$ of antipodal points on S^n we have $f(p) \neq f(-p)$.

Equivalent negation

There exists a $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map $\tilde{f}: S^n \rightarrow S^{n-1}$.

The Borsuk-Ulam Theorem

Theorem (Borsuk-Ulam)

Let n be a positive natural number. If $f: S^n \rightarrow \mathbb{R}^n$ is continuous, then there exists a pair $(p, -p)$ of antipodal points on S^n such that $f(p) = f(-p)$.

The logical negation of the theorem

There exists a continuous map $f: S^n \rightarrow \mathbb{R}^n$ such that for all pairs $(p, -p)$ of antipodal points on S^n we have $f(p) \neq f(-p)$.

Equivalent negation


There exists a $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map $\tilde{f}: S^n \rightarrow S^{n-1}$.

Theorem (equivariant formulation)

*Let n be a positive natural number. There does **not** exist a $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map $\tilde{f}: S^n \rightarrow S^{n-1}$.*


Equivariant join construction

For any topological spaces X and Y , one defines the **join** space $X * Y$ as the quotient of $[0, 1] \times X \times Y$ by a certain equivalence relation:



Equivariant join construction


For any topological spaces X and Y , one defines the **join** space $X * Y$ as the quotient of $[0, 1] \times X \times Y$ by a certain equivalence relation:



If X is a compact Hausdorff space with a continuous free action of a compact Hausdorff group G , then the diagonal action of G on the join $X * G$ is again continuous and free.

Equivariant join construction

For any topological spaces X and Y , one defines the **join** space $X * Y$ as the quotient of $[0, 1] \times X \times Y$ by a certain equivalence relation:



If X is a compact Hausdorff space with a continuous free action of a compact Hausdorff group G , then the diagonal action of G on the join $X * G$ is again continuous and free. In particular, for the antipodal action of $\mathbb{Z}/2\mathbb{Z}$ on S^{n-1} , we obtain a $\mathbb{Z}/2\mathbb{Z}$ -equivariant identification $S^n \cong S^{n-1} * \mathbb{Z}/2\mathbb{Z}$ for the antipodal and diagonal actions respectively.

Join formulation and classical generalization

Thus the Borsuk-Ulam Theorem is equivalent to:

Theorem (join formulation)

Let n be a positive natural number. There does **not** exist a $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map $\tilde{f}: S^{n-1} * \mathbb{Z}/2\mathbb{Z} \rightarrow S^{n-1}$.

Join formulation and classical generalization

Thus the Borsuk-Ulam Theorem is equivalent to:

Theorem (join formulation)

Let n be a positive natural number. There does **not** exist a $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map $\tilde{f}: S^{n-1} * \mathbb{Z}/2\mathbb{Z} \rightarrow S^{n-1}$.

This naturally leads to:

A classical Borsuk-Ulam-type conjecture

Let X be a compact Hausdorff space equipped with a continuous free action of a non-trivial compact Hausdorff group G . Then, for the diagonal action of G on $X * G$, there does **not** exist a G -equivariant continuous map $f: X * G \rightarrow X$.

Join formulation and classical generalization

Thus the Borsuk-Ulam Theorem is equivalent to:

Theorem (join formulation)

Let n be a positive natural number. There does **not** exist a $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map $\tilde{f}: S^{n-1} * \mathbb{Z}/2\mathbb{Z} \rightarrow S^{n-1}$.

This naturally leads to:

A classical Borsuk-Ulam-type conjecture

Let X be a compact Hausdorff space equipped with a continuous free action of a non-trivial compact Hausdorff group G . Then, for the diagonal action of G on $X * G$, there does **not** exist a G -equivariant continuous map $f: X * G \rightarrow X$.

At the moment, the conjecture is known to hold under the assumption of local triviality. Without this assumption, it implies a version of the Hilbert-Smith conjecture.

Research and Innovation Staff Exchange network of:
IMPAN (Poland), University of Warsaw (Poland), University of Łódź (Poland), University of Glasgow (G. Britain), University of Copenhagen (Denmark), University of Münster (Germany), Free University of Brussels (Belgium), SISSA (Italy), Penn State University (USA), University of Colorado at Boulder (USA), University of Kansas at Lawrence (USA), University of California at Berkeley (USA), University of Denver (USA), **State University of New York at Buffalo (USA)**, Fields Institute (Canada), University of New Brunswick at Fredericton (Canada), University of Wollongong (Australia), Australian National University (Australia), University Michoacana de San Nicolás de Hidalgo (Mexico).



HORIZON 2020

The EU Framework Programme for Research and Innovation

Tentative plan of conferences

New Geometry of Quantum Dynamics planned conferences:

New Geometry of Quantum Dynamics planned conferences:

- The Banach Center, Warsaw, 15 January – 19 January 2018
(approved and funded)

Tentative plan of conferences

New Geometry of Quantum Dynamics planned conferences:

- The Banach Center, Warsaw, 15 January – 19 January 2018
(approved and funded)
- The Fields Institute, Toronto, 22 July – 16 August 2019
(pending approval)

What is a compact quantum space?

Theorem (Gelfand-Naimark I)

Every *commutative C^* -algebra* is naturally isomorphic to the algebra of all continuous complex-valued vanishing-at-infinity functions on a *locally compact Hausdorff space*.

What is a compact quantum space?

Theorem (Gelfand-Naimark I)

Every *commutative C^* -algebra* is naturally isomorphic to the algebra of all continuous complex-valued vanishing-at-infinity functions on a *locally compact Hausdorff space*.

Theorem (Gelfand-Naimark II)

Every *C^* -algebra* is a complex algebra of continuous (i.e. bounded) linear operators on a complex Hilbert space that is:

What is a compact quantum space?

Theorem (Gelfand-Naimark I)

Every *commutative C^* -algebra* is naturally isomorphic to the algebra of all continuous complex-valued vanishing-at-infinity functions on a *locally compact Hausdorff space*.

Theorem (Gelfand-Naimark II)

Every *C^* -algebra* is a complex algebra of continuous (i.e. bounded) linear operators on a complex Hilbert space that is:

- 1 a topologically closed set in the norm topology of operators,

What is a compact quantum space?

Theorem (Gelfand-Naimark I)

Every *commutative C^* -algebra* is naturally isomorphic to the algebra of all continuous complex-valued vanishing-at-infinity functions on a *locally compact Hausdorff space*.

Theorem (Gelfand-Naimark II)

Every *C^* -algebra* is a complex algebra of continuous (i.e. bounded) linear operators on a complex Hilbert space that is:

- 1 a topologically closed set in the norm topology of operators,
- 2 closed under the operation of taking adjoints of operators.

What is a compact quantum space?

Theorem (Gelfand-Naimark I)

Every *commutative C*-algebra* is naturally isomorphic to the algebra of all continuous complex-valued vanishing-at-infinity functions on a *locally compact Hausdorff space*.

Theorem (Gelfand-Naimark II)

Every *C*-algebra* is a complex algebra of continuous (i.e. bounded) linear operators on a complex Hilbert space that is:

- 1 a topologically closed set in the norm topology of operators,
- 2 closed under the operation of taking adjoints of operators.

Copernican-style revolution

Given a compact Hausdorff space of points, we can define the C*-algebra of functions on the space, but the central concept is that of a commutative C*-algebras, and points appear as characters (algebra homomorphisms into \mathbb{C}) rather than as primary objects. We think of noncommutative unital C*-algebras as algebras of functions on *compact quantum spaces*.

What is a compact quantum group?

Definition (S. L. Woronowicz)

A **compact quantum group** is a unital C^* -algebra H with a given unital $*$ -homomorphism $\Delta: H \rightarrow H \otimes_{\min} H$ such that the diagram

$$\begin{array}{ccc}
 H & \xrightarrow{\Delta} & H \otimes_{\min} H \\
 \downarrow \Delta & & \downarrow \Delta \otimes \text{id} \\
 H \otimes_{\min} H & \xrightarrow{\text{id} \otimes \Delta} & H \otimes_{\min} H \otimes_{\min} H
 \end{array}$$

commutes and the two-sided cancellation property holds:

$$\{(a \otimes 1)\Delta(b) \mid a, b \in H\}^{\text{cls}} = H \otimes_{\min} H = \{\Delta(a)(1 \otimes b) \mid a, b \in H\}^{\text{cls}}.$$

Here “cls” stands for “closed linear span”.

Free actions of compact quantum groups

Let A be a unital C^* -algebra and $\delta : A \rightarrow A \otimes_{\min} H$ a unital $*$ -homomorphism. We call δ a **coaction** (or an action of the compact quantum group (H, Δ) on A) iff

- 1 $(\delta \otimes \text{id}) \circ \delta = (\text{id} \otimes \Delta) \circ \delta$ (coassociativity),
- 2 $\{\delta(a)(1 \otimes h) \mid a \in A, h \in H\}^{\text{cls}} = A \otimes_{\min} H$ (counitality),
- 3 $\ker \delta = 0$ (injectivity).

Free actions of compact quantum groups

Let A be a unital C^* -algebra and $\delta : A \rightarrow A \otimes_{\min} H$ a unital $*$ -homomorphism. We call δ a **coaction** (or an action of the compact quantum group (H, Δ) on A) iff

- 1 $(\delta \otimes \text{id}) \circ \delta = (\text{id} \otimes \Delta) \circ \delta$ (coassociativity),
- 2 $\{\delta(a)(1 \otimes h) \mid a \in A, h \in H\}^{\text{cls}} = A \otimes_{\min} H$ (counitality),
- 3 $\ker \delta = 0$ (injectivity).

Definition (D. A. Ellwood)

A coaction δ is called **free** iff

$$\{(x \otimes 1)\delta(y) \mid x, y \in A\}^{\text{cls}} = A \otimes_{\min} H .$$

Equivariant noncommutative join construction

Definition (L. Dąbrowski, T. Hadfield, P. M. H.)

For any compact quantum group (H, Δ) acting freely on a unital C^* -algebra A , we define its **equivariant join** with H to be the unital C^* -algebra

$$A \overset{\delta}{\circledast} H := \left\{ f \in C([0, 1], A) \underset{\min}{\otimes} H \mid f(0) \in \mathbb{C} \otimes H, f(1) \in \delta(A) \right\}.$$

Equivariant noncommutative join construction

Definition (L. Dąbrowski, T. Hadfield, P. M. H.)

For any compact quantum group (H, Δ) acting freely on a unital C^* -algebra A , we define its **equivariant join** with H to be the unital C^* -algebra

$$A \overset{\delta}{\ast} H := \left\{ f \in C([0, 1], A) \underset{\min}{\otimes} H \mid f(0) \in \mathbb{C} \otimes H, f(1) \in \delta(A) \right\}.$$

Theorem (P. F. Baum, K. De Commer, P. M. H.)

The \ast -homomorphism

$$\text{id} \otimes \Delta: C([0, 1], A) \underset{\min}{\otimes} H \longrightarrow C([0, 1], A) \underset{\min}{\otimes} H \underset{\min}{\otimes} H$$

defines a free action of the compact quantum group (H, Δ) on the equivariant join C^ -algebra $A \overset{\delta}{\ast} H$.*

Noncommutative Borsuk-Ulam-type conjectures

Conjecture 1

Let A be a unital C^* -algebra with a free action of a non-trivial compact quantum group (H, Δ) . Then there **does not exist an H -equivariant $*$ -homomorphism $A \rightarrow A \otimes^{\delta} H$** . (Known to hold for (H, Δ) with classical torsion.)

Noncommutative Borsuk-Ulam-type conjectures

Conjecture 1

Let A be a unital C^* -algebra with a free action of a non-trivial compact quantum group (H, Δ) . Then there **does not exist an H -equivariant $*$ -homomorphism $A \rightarrow A \otimes^{\delta} H$** . (Known to hold for (H, Δ) with classical torsion.)

Conjecture 2

Let A be a unital C^* -algebra with a free action of a non-trivial compact quantum group (H, Δ) . If A admits a character, then there **does not exist an H -equivariant $*$ -homomorphism $H \rightarrow A \otimes^{\delta} H$** . (Follows from Conjecture 1.)

Noncommutative Borsuk-Ulam-type conjectures

Conjecture 1

Let A be a unital C^* -algebra with a free action of a non-trivial compact quantum group (H, Δ) . Then there **does not exist an H -equivariant $*$ -homomorphism $A \rightarrow A \otimes^{\delta} H$** . (Known to hold for (H, Δ) with classical torsion.)

Conjecture 2

Let A be a unital C^* -algebra with a free action of a non-trivial compact quantum group (H, Δ) . If A admits a character, then there **does not exist an H -equivariant $*$ -homomorphism $H \rightarrow A \otimes^{\delta} H$** . (Follows from Conjecture 1.)

Classical cases

If X is a compact Hausdorff principal G -bundle, $A = C(X)$ and $H = C(G)$, then Conjecture 2 states that the principal G -bundle $X * G$ is not trivializable unless G is trivial. This is clearly true because otherwise **$G * G$ would be trivializable, which is tantamount to G being contractible**, and the only contractible compact Hausdorff group is the trivial one.

Iterated joins of the quantum $SU(2)$ group

Consider the fibration defining the quaternionic projective space:

$$SU(2) * \cdots * SU(2) \cong S^{4n+3}, \quad S^{4n+3}/SU(2) = \mathbb{H}P^n.$$

To obtain a q -deformation of this fibration, we take $H := C(SU_q(2))$ and $A := C(S_q^{4n+3})$ equal to the n -times iterated equivariant join of H . The thus given quantum principal $SU_q(2)$ -bundle is *not* trivializable:

Theorem

There does *not* exist a $C(SU_q(2))$ -equivariant $*$ -homomorphism

$$f: C(SU_q(2)) \longrightarrow C(S_q^{4n+3}) \otimes^\delta C(SU_q(2)).$$

Iterated joins of the quantum $SU(2)$ group

Consider the fibration defining the quaternionic projective space:

$$SU(2) * \cdots * SU(2) \cong S^{4n+3}, \quad S^{4n+3}/SU(2) = \mathbb{H}P^n.$$

To obtain a q -deformation of this fibration, we take $H := C(SU_q(2))$ and $A := C(S_q^{4n+3})$ equal to the n -times iterated equivariant join of H . The thus given quantum principal $SU_q(2)$ -bundle is *not* trivializable:

Theorem

There does *not* exist a $C(SU_q(2))$ -equivariant $*$ -homomorphism

$$f: C(SU_q(2)) \longrightarrow C(S_q^{4n+3}) \otimes^\delta C(SU_q(2)).$$

This theorem holds because $SU_q(2)$ has classical torsion elements. It also follows from the stable non-triviality of the dual tautological quaternionic line bundle:

The tautological quaternionic line bundle

If f existed, there would exist an equivariant map F

$$C(SU_q(2)) \rightarrow C(S_q^{4n+3}) \otimes^{\delta} C(SU_q(2)) \rightarrow C(SU_q(2)) \otimes^{\Delta} C(SU_q(2)).$$

The tautological quaternionic line bundle

If f existed, there would exist an equivariant map F

$$C(SU_q(2)) \rightarrow C(S_q^{4n+3}) \otimes^{\delta} C(SU_q(2)) \rightarrow C(SU_q(2)) \otimes^{\Delta} C(SU_q(2)).$$

Furthermore, for any finite-dimensional representation V of a compact quantum group (H, Δ) , the associated finitely generated projective module $(H \otimes^{\Delta} H) \square_H V$ is represented by a Milnor idempotent $p_{U^{-1}}$, where U is a matrix of the representation V , and an even index pairing calculation for $p_{U^{-1}}$ might be replaced by an odd index pairing calculation for U .

The tautological quaternionic line bundle

If f existed, there would exist an equivariant map F

$$C(SU_q(2)) \rightarrow C(S_q^{4n+3}) \otimes^{\delta} C(SU_q(2)) \rightarrow C(SU_q(2)) \otimes^{\Delta} C(SU_q(2)).$$

Furthermore, for any finite-dimensional representation V of a compact quantum group (H, Δ) , the associated finitely generated projective module $(H \otimes^{\Delta} H) \square_H V$ is represented by a Milnor idempotent $p_{U^{-1}}$, where U is a matrix of the representation V , and an even index pairing calculation for $p_{U^{-1}}$ might be replaced by an odd index pairing calculation for U .

Now, for $H := C(SU_q(2))$ and V the fundamental representation of $SU_q(2)$, the module $(H \otimes^{\Delta} H) \square_H V$ is the section module of the **dual tautological quaternionic line bundle**. It is *not* stably free by the non-vanishing of an index pairing of the fundamental representation of $SU_q(2)$ with an appropriate odd Fredholm module. This contradicts the existence of F .