



# THERE AND BACK AGAIN: FROM THE BORSUK-ULAM THEOREM TO QUANTUM SPACES

**Piotr M. Hajac** (IMPAN / University of New Brunswick)  
**Mariusz Tobolski** (Warsaw University)

Joint work with  
Paul F. Baum, Ludwik Dąbrowski and Tomasz Maszczyk

Zakopane, 9 February 2016

1996 m



1996 m



# A mystery hiker at 1996 m





Jiří Matoušek

# Using the Borsuk-Ulam Theorem

Lectures on Topological Methods  
in Combinatorics and Geometry



# The Borsuk-Ulam Theorem

## Theorem (Borsuk-Ulam)

*Let  $n$  be a positive natural number. If  $f: S^n \rightarrow \mathbb{R}^n$  is continuous, then there exists a pair  $(p, -p)$  of antipodal points on  $S^n$  such that  $f(p) = f(-p)$ .*

# The Borsuk-Ulam Theorem

## Theorem (Borsuk-Ulam)

*Let  $n$  be a positive natural number. If  $f: S^n \rightarrow \mathbb{R}^n$  is continuous, then there exists a pair  $(p, -p)$  of antipodal points on  $S^n$  such that  $f(p) = f(-p)$ .*

## The logical negation of the theorem

There exists a continuous map  $f: S^n \rightarrow \mathbb{R}^n$  such that for all pairs  $(p, -p)$  of antipodal points on  $S^n$  we have  $f(p) \neq f(-p)$ .

# The Borsuk-Ulam Theorem

## Theorem (Borsuk-Ulam)

*Let  $n$  be a positive natural number. If  $f: S^n \rightarrow \mathbb{R}^n$  is continuous, then there exists a pair  $(p, -p)$  of antipodal points on  $S^n$  such that  $f(p) = f(-p)$ .*

## The logical negation of the theorem

There exists a continuous map  $f: S^n \rightarrow \mathbb{R}^n$  such that for all pairs  $(p, -p)$  of antipodal points on  $S^n$  we have  $f(p) \neq f(-p)$ .

## Equivalent negation

There exists a  $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map  $\tilde{f}: S^n \rightarrow S^{n-1}$ .

# The Borsuk-Ulam Theorem

## Theorem (Borsuk-Ulam)

*Let  $n$  be a positive natural number. If  $f: S^n \rightarrow \mathbb{R}^n$  is continuous, then there exists a pair  $(p, -p)$  of antipodal points on  $S^n$  such that  $f(p) = f(-p)$ .*

## The logical negation of the theorem

There exists a continuous map  $f: S^n \rightarrow \mathbb{R}^n$  such that for all pairs  $(p, -p)$  of antipodal points on  $S^n$  we have  $f(p) \neq f(-p)$ .

## Equivalent negation

There exists a  $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map  $\tilde{f}: S^n \rightarrow S^{n-1}$ .


## Theorem (equivariant formulation)

*Let  $n$  be a positive natural number. There does **not** exist a  $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map  $\tilde{f}: S^n \rightarrow S^{n-1}$ .*




# Equivariant join construction

For any topological spaces  $X$  and  $Y$ , one defines the **join** space  $X * Y$  as the quotient of  $[0, 1] \times X \times Y$  by a certain equivalence relation:



## Equivariant join construction


For any topological spaces  $X$  and  $Y$ , one defines the **join** space  $X * Y$  as the quotient of  $[0, 1] \times X \times Y$  by a certain equivalence relation:



If  $X$  is a compact Hausdorff space with a continuous free action of a compact Hausdorff group  $G$ , then the diagonal action of  $G$  on the join  $X * G$  is again continuous and free.

# Equivariant join construction

For any topological spaces  $X$  and  $Y$ , one defines the **join** space  $X * Y$  as the quotient of  $[0, 1] \times X \times Y$  by a certain equivalence relation:



If  $X$  is a compact Hausdorff space with a continuous free action of a compact Hausdorff group  $G$ , then the diagonal action of  $G$  on the join  $X * G$  is again continuous and free. In particular, for the antipodal action of  $\mathbb{Z}/2\mathbb{Z}$  on  $S^{n-1}$ , we obtain a  $\mathbb{Z}/2\mathbb{Z}$ -equivariant identification  $S^n \cong S^{n-1} * \mathbb{Z}/2\mathbb{Z}$  for the antipodal and diagonal actions respectively.

# Join formulation and classical generalization

Thus the Borsuk-Ulam Theorem is equivalent to:

Theorem (join formulation)

*Let  $n$  be a positive natural number. There does **not** exist a  $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map  $\tilde{f}: S^{n-1} * \mathbb{Z}/2\mathbb{Z} \rightarrow S^{n-1}$ .*

# Join formulation and classical generalization

Thus the Borsuk-Ulam Theorem is equivalent to:

## Theorem (join formulation)

Let  $n$  be a positive natural number. There does **not** exist a  $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map  $\tilde{f}: S^{n-1} * \mathbb{Z}/2\mathbb{Z} \rightarrow S^{n-1}$ .

This naturally leads to:

## A classical Borsuk-Ulam-type conjecture

Let  $X$  be a compact Hausdorff space equipped with a continuous free action of a non-trivial compact Hausdorff group  $G$ . Then, for the diagonal action of  $G$  on  $X * G$ , there does **not** exist a  $G$ -equivariant continuous map  $f: X * G \rightarrow X$ .



# Join formulation and classical generalization

Thus the Borsuk-Ulam Theorem is equivalent to:

## Theorem (join formulation)

Let  $n$  be a positive natural number. There does **not** exist a  $\mathbb{Z}/2\mathbb{Z}$ -equivariant continuous map  $\tilde{f}: S^{n-1} * \mathbb{Z}/2\mathbb{Z} \rightarrow S^{n-1}$ .


This naturally leads to:

## A classical Borsuk-Ulam-type conjecture

Let  $X$  be a compact Hausdorff space equipped with a continuous free action of a non-trivial compact Hausdorff group  $G$ . Then, for the diagonal action of  $G$  on  $X * G$ , there does **not** exist a  $G$ -equivariant continuous map  $f: X * G \rightarrow X$ .

## Corollary

*Ageev's conjecture about the Menger compacta.*




SIMONS  
FOUNDATION

1 Sep – 30 Nov 2016, Simons Semester in the Banach Center

**NONCOMMUTATIVE GEOMETRY THE NEXT GENERATION**

*Paul F. Baum, Alan Carey, Piotr M. Hajac, Tomasz Maszczyk*




SIMONS  
FOUNDATION

1 Sep – 30 Nov 2016, Simons Semester in the Banach Center

## **NONCOMMUTATIVE GEOMETRY THE NEXT GENERATION**

*Paul F. Baum, Alan Carey, Piotr M. Hajac, Tomasz Maszczyk*

Funding available for longer stays (Senior Professors and Junior Professors, Postdocs, or PhD Students).



**SIMONS**  
FOUNDATION

1 Sep – 30 Nov 2016, Simons Semester in the Banach Center

## **NONCOMMUTATIVE GEOMETRY THE NEXT GENERATION**

*Paul F. Baum, Alan Carey, Piotr M. Hajac, Tomasz Maszczyk*

Funding available for longer stays (Senior Professors and Junior Professors, Postdocs, or PhD Students).



# Noncommutative Geometry the Next Generation

- ① 4–17 September, Będlewo & Warsaw, school on  
Noncommutative geometry and quantum groups



# Noncommutative Geometry the Next Generation

- ① 4–17 September, Będlewo & Warsaw, school on  
Noncommutative geometry and quantum groups
- ② 19 September – 14 October, 20-hour lecture course  
An invitation to  $C^*$ -algebras by Karen R. Strung

# Noncommutative Geometry the Next Generation

- ① 4–17 September, Będlewo & Warsaw, school on  
Noncommutative geometry and quantum groups
- ② 19 September – 14 October, 20-hour lecture course  
An invitation to  $C^*$ -algebras by Karen R. Strung
- ③ 19 September – 14 October, 20-hour lecture course  
An invitation to Hopf algebras by Réamonn Ó Buachalla

# Noncommutative Geometry the Next Generation

- ① 4–17 September, Będlewo & Warsaw, school on  
**Noncommutative geometry and quantum groups**
- ② 19 September – 14 October, 20-hour lecture course  
**An invitation to  $C^*$ -algebras** by Karen R. Strung
- ③ 19 September – 14 October, 20-hour lecture course  
**An invitation to Hopf algebras** by Réamonn Ó Buachalla
- ④ 19 September – 14 October, 20-hour lecture course  
**Noncommutative topology for beginners** by Tatiana Shulman

# Noncommutative Geometry the Next Generation

- ① 4–17 September, Będlewo & Warsaw, school on  
**Noncommutative geometry and quantum groups**
- ② 19 September – 14 October, 20-hour lecture course  
**An invitation to  $C^*$ -algebras** by Karen R. Strung
- ③ 19 September – 14 October, 20-hour lecture course  
**An invitation to Hopf algebras** by Réamonn Ó Buachalla
- ④ 19 September – 14 October, 20-hour lecture course  
**Noncommutative topology for beginners** by Tatiana Shulman
- ⑤ 17–21 Oct. **Cyclic homology**  
J. Cuntz, P. M. Hajac, T. Maszczyk, R. Nest

# Noncommutative Geometry the Next Generation

- ① 4–17 September, Będlewo & Warsaw, school on  
**Noncommutative geometry and quantum groups**
- ② 19 September – 14 October, 20-hour lecture course  
**An invitation to  $C^*$ -algebras** by Karen R. Strung
- ③ 19 September – 14 October, 20-hour lecture course  
**An invitation to Hopf algebras** by Réamonn Ó Buachalla
- ④ 19 September – 14 October, 20-hour lecture course  
**Noncommutative topology for beginners** by Tatiana Shulman
- ⑤ 17–21 Oct. **Cyclic homology**  
J. Cuntz, P. M. Hajac, T. Maszczyk, R. Nest
- ⑥ 24–28 Oct. **Noncommutative index theory**  
P. F. Baum, A. Carey, M. J. Pflaum, A. Sitarz

# Noncommutative Geometry the Next Generation

- ① 4–17 September, Będlewo & Warsaw, school on  
**Noncommutative geometry and quantum groups**
- ② 19 September – 14 October, 20-hour lecture course  
**An invitation to  $C^*$ -algebras** by Karen R. Strung
- ③ 19 September – 14 October, 20-hour lecture course  
**An invitation to Hopf algebras** by Réamonn Ó Buachalla
- ④ 19 September – 14 October, 20-hour lecture course  
**Noncommutative topology for beginners** by Tatiana Shulman
- ⑤ 17–21 Oct. **Cyclic homology**  
J. Cuntz, P. M. Hajac, T. Maszczyk, R. Nest
- ⑥ 24–28 Oct. **Noncommutative index theory**  
P. F. Baum, A. Carey, M. J. Pflaum, A. Sitarz
- ⑦ 14–18 Nov. **Topological quantum groups and Hopf algebras**  
K. De Commer, P. M. Hajac, R. Ó Buachalla, A. Skalski

# Noncommutative Geometry the Next Generation

- ① 4–17 September, Będlewo & Warsaw, school on  
**Noncommutative geometry and quantum groups**
- ② 19 September – 14 October, 20-hour lecture course  
**An invitation to  $C^*$ -algebras** by Karen R. Strung
- ③ 19 September – 14 October, 20-hour lecture course  
**An invitation to Hopf algebras** by Réamonn Ó Buachalla
- ④ 19 September – 14 October, 20-hour lecture course  
**Noncommutative topology for beginners** by Tatiana Shulman
- ⑤ 17–21 Oct. **Cyclic homology**  
J. Cuntz, P. M. Hajac, T. Maszczyk, R. Nest
- ⑥ 24–28 Oct. **Noncommutative index theory**  
P. F. Baum, A. Carey, M. J. Pflaum, A. Sitarz
- ⑦ 14–18 Nov. **Topological quantum groups and Hopf algebras**  
K. De Commer, P. M. Hajac, R. Ó Buachalla, A. Skalski
- ⑧ 21–25 Nov. **Structure and classification of  $C^*$ -algebras**  
G. Elliott, K. R. Strung, W. Winter, J. Zacharias

**GEOMETRY, REPRESENTATION THEORY  
AND THE BAUM-CONNES CONJECTURE**

A workshop in honour of **Paul F. Baum** on the occasion of his 80th birthday organized by Alan Carey, George Elliott, Piotr M. Hajac, and Ryszard Nest.



## GEOMETRY, REPRESENTATION THEORY AND THE BAUM-CONNES CONJECTURE

A workshop in honour of **Paul F. Baum** on the occasion of his 80th birthday organized by Alan Carey, George Elliott, Piotr M. Hajac, and Ryszard Nest.

Sponsored by:

- The Fields Institute, University of Toronto, Canada
- National Science Foundation, USA
- The Pennsylvania State University, USA



FIELDS



# Free actions of compact quantum groups

Let  $A$  be a unital  $C^*$ -algebra and  $\delta : A \rightarrow A \otimes_{\min} H$  a unital  $*$ -homomorphism. We call  $\delta$  a **coaction** of  $H$  on  $A$  (or an action of the compact quantum group  $(H, \Delta)$  on  $A$ ) iff

- 1  $(\delta \otimes \text{id}) \circ \delta = (\text{id} \otimes \Delta) \circ \delta$  (coassociativity),
- 2  $\{\delta(a)(1 \otimes h) \mid a \in A, h \in H\}^{\text{cls}} = A \otimes_{\min} H$  (counitality)
- 3  $\ker \delta = 0$  (injectivity).

# Free actions of compact quantum groups

Let  $A$  be a unital  $C^*$ -algebra and  $\delta : A \rightarrow A \otimes_{\min} H$  a unital  $*$ -homomorphism. We call  $\delta$  a **coaction** of  $H$  on  $A$  (or an action of the compact quantum group  $(H, \Delta)$  on  $A$ ) iff

- 1  $(\delta \otimes \text{id}) \circ \delta = (\text{id} \otimes \Delta) \circ \delta$  (coassociativity),
- 2  $\{\delta(a)(1 \otimes h) \mid a \in A, h \in H\}^{\text{cls}} = A \otimes_{\min} H$  (counitality)
- 3  $\ker \delta = 0$  (injectivity).

Definition (D. A. Ellwood)

A coaction  $\delta$  is called **free** iff

$$\{(x \otimes 1)\delta(y) \mid x, y \in A\}^{\text{cls}} = A \otimes_{\min} H .$$

# Equivariant noncommutative join construction

Definition (L. Dąbrowski, T. Hadfield, P. M. H.)

For any compact quantum group  $(H, \Delta)$  acting freely on a unital  $C^*$ -algebra  $A$ , we define its **equivariant join** with  $H$  to be the unital  $C^*$ -algebra

$$A \overset{\delta}{\circledast} H := \left\{ f \in C([0, 1], A) \underset{\min}{\otimes} H \mid f(0) \in \mathbb{C} \otimes H, f(1) \in \delta(A) \right\}.$$

# Equivariant noncommutative join construction

Definition (L. Dąbrowski, T. Hadfield, P. M. H.)

For any compact quantum group  $(H, \Delta)$  acting freely on a unital  $C^*$ -algebra  $A$ , we define its **equivariant join** with  $H$  to be the unital  $C^*$ -algebra

$$A \overset{\delta}{\ast} H := \left\{ f \in C([0, 1], A) \underset{\min}{\otimes} H \mid f(0) \in \mathbb{C} \otimes H, f(1) \in \delta(A) \right\}.$$

Theorem (P. F. Baum, K. De Commer, P. M. H.)

*The  $\ast$ -homomorphism*

$$\text{id} \otimes \Delta: C([0, 1], A) \underset{\min}{\otimes} H \longrightarrow C([0, 1], A) \underset{\min}{\otimes} H \underset{\min}{\otimes} H$$

*defines a free action of the compact quantum group  $(H, \Delta)$  on the equivariant join  $C^*$ -algebra  $A \overset{\delta}{\ast} H$ .*

# Noncommutative Borsuk-Ulam-type conjectures

## Conjecture 1

Let  $A$  be a unital  $C^*$ -algebra with a free action of a non-trivial compact quantum group  $(H, \Delta)$ . Then there **does not exist an  $H$ -equivariant  $*$ -homomorphism  $A \rightarrow A \otimes^{\delta} H$ .**

# Noncommutative Borsuk-Ulam-type conjectures

## Conjecture 1

Let  $A$  be a unital  $C^*$ -algebra with a free action of a non-trivial compact quantum group  $(H, \Delta)$ . Then there **does not exist an  $H$ -equivariant  $*$ -homomorphism  $A \rightarrow A \otimes^{\delta} H$ .**

## Conjecture 2

Let  $A$  be a unital  $C^*$ -algebra with a free action of a non-trivial compact quantum group  $(H, \Delta)$ . Then there **does not exist an  $H$ -equivariant  $*$ -homomorphism  $H \rightarrow A \otimes^{\delta} H$ .**

# Noncommutative Borsuk-Ulam-type conjectures

## Conjecture 1

Let  $A$  be a unital  $C^*$ -algebra with a free action of a non-trivial compact quantum group  $(H, \Delta)$ . Then there **does not exist an  $H$ -equivariant  $*$ -homomorphism  $A \rightarrow A \otimes^{\delta} H$** .

## Conjecture 2

Let  $A$  be a unital  $C^*$ -algebra with a free action of a non-trivial compact quantum group  $(H, \Delta)$ . Then there **does not exist an  $H$ -equivariant  $*$ -homomorphism  $H \rightarrow A \otimes^{\delta} H$** .

## The classical case

If  $X$  is a compact Hausdorff principal  $G$ -bundle,  $A = C(X)$  and  $H = C(G)$ , then Conjecture 2 states that the principal  $G$ -bundle  $X * G$  is not trivializable unless  $G$  is trivial. This is clearly true because otherwise  **$G * G$  would be trivializable, which is tantamount to  $G$  being contractible**, and the only contractible compact Hausdorff group is trivial.



# Noncommutative Borsuk-Ulam-type conjectures

## Conjecture 1

Let  $A$  be a unital  $C^*$ -algebra with a free action of a non-trivial compact quantum group  $(H, \Delta)$ . Then there **does not exist an  $H$ -equivariant  $*$ -homomorphism  $A \rightarrow A \otimes^{\delta} H$** .

## Conjecture 2

Let  $A$  be a unital  $C^*$ -algebra with a free action of a non-trivial compact quantum group  $(H, \Delta)$ . Then there **does not exist an  $H$ -equivariant  $*$ -homomorphism  $H \rightarrow A \otimes^{\delta} H$** .

## The classical case

If  $X$  is a compact Hausdorff principal  $G$ -bundle,  $A = C(X)$  and  $H = C(G)$ , then Conjecture 2 states that the principal  $G$ -bundle  $X * G$  is not trivializable unless  $G$  is trivial. This is clearly true because otherwise  **$G * G$  would be trivializable, which is tantamount to  $G$  being contractible**, and the only contractible compact Hausdorff group is trivial. On the other hand, we do not know whether Conjecture 1 holds in general in the commutative case.

## Iterated joins of the quantum $SU(2)$ group

Consider the defining fibration of the quaternionic projective space:

$$SU(2) * \cdots * SU(2) \cong S^{4n+3}, \quad S^{4n+3}/SU(2) = \mathbb{H}P^n.$$

To obtain a  $q$ -deformation of this fibration, we take  $H = C(SU_q(2))$  and  $A$  equal to a finitely iterated equivariant join of  $H$ . The quantum principal  $SU_q(2)$ -bundle thus given is *not* trivializable:

### Theorem (main)

There does *not* exist a  $C(SU_q(2))$ -equivariant  $*$ -homomorphism  $f: C(SU_q(2)) \rightarrow A \otimes^\delta C(SU_q(2))$ .

# Iterated joins of the quantum $SU(2)$ group

Consider the defining fibration of the quaternionic projective space:

$$SU(2) * \cdots * SU(2) \cong S^{4n+3}, \quad S^{4n+3}/SU(2) = \mathbb{H}P^n.$$

To obtain a  $q$ -deformation of this fibration, we take  $H = C(SU_q(2))$  and  $A$  equal to a finitely iterated equivariant join of  $H$ . The quantum principal  $SU_q(2)$ -bundle thus given is *not* trivializable:

## Theorem (main)

There does *not* exist a  $C(SU_q(2))$ -equivariant  $*$ -homomorphism  $f: C(SU_q(2)) \rightarrow A \otimes^\delta C(SU_q(2))$ .

**Proof outline:** First, we prove that for any finite-dimensional representation  $V$  of a compact quantum group  $(H, \Delta)$ , the associated finitely-generated projective module  $(H \otimes^\Delta H) \square_H V$  is represented by a Milnor idempotent  $p_{U^{-1}}$ , where  $U$  is a matrix of the representation  $V$ . Then we choose  $V$  to be the fundamental representation of  $SU_q(2)$ , and infer from index pairing considerations that  $(H \otimes^\Delta H) \square_H V$  is not stably free. Finally, by the Pulling Back Theorem, we conclude that  $(A \otimes^\delta H) \square_H V$  is *not* stably free, whence  $f$  does not exist. □

**Research and Innovation Staff Exchange** network of:  
IMPAN (Poland), **University of Warsaw** (Poland), University of Łódź (Poland), University of Glasgow (G. Britain), University of Aberdeen (G. Britain), University of Copenhagen (Denmark), University of Münster (Germany), Free University of Brussels (Belgium), SISSA (Italy), Penn State University (USA), University of Colorado at Boulder (USA), University of Kansas at Lawrence (USA), University of California at Berkeley (USA), University of Denver (USA), Fields Institute (Canada), University of New Brunswick at Fredericton (Canada), University of Wollongong (Australia), Australian National University (Australia), University of Otago (New Zealand), University Michoacana de San Nicolás de Hidalgo (Mexico).



**HORIZON 2020**

The EU Framework Programme for Research and Innovation